

## MaineHealth MaineHealth Knowledge Connection

Maine Medical Center

All MaineHealth

5-1-2019

# Feasibility of pre-operative mTOR inhibitor Sirolimus in children and young adults with desmoid tumor

Stephanie Verwys  
*Maine Medical Center*

Clara Magyar

Kathleen Glick  
*Maine Medical Center*

Douglas Hawkins

Archana Sharma

*See next page for additional authors*

Follow this and additional works at: <https://knowledgeconnection.mainehealth.org/mmc>

 Part of the [Pediatrics Commons](#), and the [Surgery Commons](#)

### Recommended Citation

Verwys, Stephanie; Magyar, Clara; Glick, Kathleen; Hawkins, Douglas; Sharma, Archana; Weigel, Brenda; Chastain, Katherine; Khoury, Joseph; Manalang, Michele; Dry, Sarah; Federman, Noah; and Weiss, Aaron, "Feasibility of pre-operative mTOR inhibitor Sirolimus in children and young adults with desmoid tumor" (2019). *Maine Medical Center*. 700.  
<https://knowledgeconnection.mainehealth.org/mmc/700>

This Poster is brought to you for free and open access by the All MaineHealth at MaineHealth Knowledge Connection. It has been accepted for inclusion in Maine Medical Center by an authorized administrator of MaineHealth Knowledge Connection. For more information, please contact [mckeld1@mmc.org](mailto:mckeld1@mmc.org).

---

**Authors**

Stephanie Verwys, Clara Magyar, Kathleen Glick, Douglas Hawkins, Archana Sharma, Brenda Weigel, Katherine Chastain, Joseph Khoury, Michele Manalang, Sarah Dry, Noah Federman, and Aaron Weiss



# Feasibility of pre-operative mTOR inhibitor Sirolimus in children and young adults with desmoid tumor

Stephanie Verwys<sup>1</sup>, Clara Magyar<sup>2</sup>, Kathleen Glick<sup>1</sup>, Douglas Hawkins<sup>3</sup>, Archana Sharma<sup>4</sup>, Brenda Weigel<sup>5</sup>, Katherine Chastain<sup>6</sup>, Joseph Khoury<sup>7</sup>, Michele Manalang<sup>8</sup>, Sarah Dry<sup>2</sup>, Noah Federman<sup>2</sup>, and Aaron Weiss<sup>1</sup>

<sup>1</sup>Maine Medical Center, Portland, ME, <sup>2</sup>The David Geffen School of Medicine at UCLA, Los Angeles, CA, <sup>3</sup>Seattle Children's Hospital, Seattle, WA, <sup>4</sup>Rutgers Cancer Institute of New Jersey, New Brunswick, NJ, <sup>5</sup>University of Minnesota, Masonic Cancer Center, Minneapolis, MN, <sup>6</sup>Mercy Hospital, University of Missouri-Kansas City, Kansas City, MO, <sup>7</sup>The University of Texas, MD Anderson Cancer Center, Houston, TX, <sup>8</sup>Marshfield Clinic, Marshfield, WI

## Background

- Desmoid tumor represents an intermediate grade neoplasm with a striking predilection for locally invasive growth and recurrence following resection
- More effective, well-tolerated non-surgical treatment options are needed
- Current approaches**
  - If feasible, watchful waiting is the preferred approach
    - 20-30% spontaneous regression
  - In situations where treatment is indicated, the following approaches are utilized
    - Surgery is the primary approach if minimal morbidity is anticipated
    - Medical therapies
      - Cytotoxic drugs
      - Tyrosine kinase inhibitors
      - Hydroxyurea
      - Gamma secretase inhibitors
- mTOR Inhibitor Rationale**
  - Desmoid tumor is well-known to be associated with deregulation of the APC/β-catenin pathway
  - Deregulation of the mTOR cell proliferation/survival pathway may play an important role in tumor biology when the APC/β-catenin pathway is disrupted
  - The mTOR inhibitor **sirolimus** is attractive as a potential targeted therapy for desmoid tumor
    - Well-tolerated in children and young adults
    - Can be given orally in tablet or liquid formulation


## Objectives

- Primary**
  - To determine whether mTOR pathway activation decreases in patients with surgically resectable desmoid tumor that is removed following pre-operative treatment with sirolimus
- Secondary**
  - To assess whether sirolimus improves desmoid tumor-associated pain
  - To begin to explore whether pre-operative sirolimus decreases tumor recurrence following surgical removal of desmoid tumor felt to be at high-risk for recurrence because of size and/or anatomic site
  - To assess the safety and tolerability of pre-operative sirolimus in patients with desmoid tumor

## Methods

- Multi-institutional study open and actively accruing patients
- Eligibility criteria
  - <30 years of age
  - Surgery is planned to remove their desmoid tumor and either
    - (a) the desmoid tumor has already recurred after a prior surgery or
    - (b) the newly diagnosed or previously unresected disease is judged to be at high risk for recurrence due to its size (>5 cm) or location at an anatomic site making it unlikely to be resected with negative margins (e.g., adjacent to neurovascular structures)
  - Patients with germline APC causing FAP/Gardner's syndrome
- This is an IRB-approved study and patient consent is required

Figure 1. Experimental Design Schema




## Results

- Nine of an anticipated 15 total patients have enrolled to date
- Ages have ranged from 5 to 28 years
- All patients have been able to take the pre-operative sirolimus as prescribed and undergone surgery within the protocol-directed time frame
- All toxicities have been as expected and Common Terminology Criteria for Adverse Events grade 1 and 2 only except for one grade 3 neutropenia
- No post-operative complications have been reported
- IHC staining is ongoing for p4E-BP1, p70S6K, and pAKT (Figure 2)
- Pain assessment measurements and anatomical imaging are being performed at designated surveillance intervals

## Histologic Assessment

- Tissue
  - Pre-therapy biopsy
  - Post-therapy biopsy
  - Archived specimens (future analysis)
    - Paired, pre-treatment
    - Non-chemotherapy treatment

Figure 2. Representative pre- and post-operative IHC staining for mTOR pathway proteins p4E-BP1, p70S6K, and pAKT in desmoid tumor



## Conclusion

- Sirolimus appears to be well-tolerated when administered in the pre-operative setting to children and young adults with desmoid tumor
- Surgery is feasible and safe immediately after completing therapy
- Formal assessment of the mTOR pathway by IHC analysis will take place at study completion
- The study continues to actively accrue

## Acknowledgements

Thank you to the Desmoid Tumor Research Foundation for funding, Pfizer for drug supply, and all of the participating sites, and patients.